## AST 250 – Spring 2019 Homework Due: Monday Feb. 11th

14. In this problem, you will derive the gravitational potential energy,  $U_g$ , of a uniform density sphere ( $\rho$  = constant). Consider the geometry of a test mass, dm<sub>i</sub>, within a thin spherical shell with differential radius dr:



The test mass has a gravitational potential energy of:

$$dU_{g,i}=-Grac{M_r\,dm_i}{r}$$

where  $M_r$  is the total mass interior to radius r. The gravitational potential energy of the shell is then found by substituting dm = volume of shell \* density =  $4\pi r^2 dr * \rho$  to obtain

$$dU_g = -Grac{M_r\,4\pi r^2
ho}{r}\,dr$$

(a) To calculate the total gravitational potential energy of a sphere Ug, we need to integrate dUg over all mass shells from the center to total radius R (integrating over dr). Write down this integral and pull everything that doesn't depend on r outside the integral. NOTE: *capital* R is the total radius of the sphere. Little r is a variable than goes from 0 to R.

(b) How is  $M_r$  related to r and  $\rho$ ? Assume  $\rho$  is constant. Substitute for  $M_r$  and evaluate the integral. Your answer should only contain numbers, G, and  $\rho$  and R as variables.

(c) Now substitute for  $\rho$  to convert you answer to only have numbers, G, and M (total mass) and R as variables. This is a good equation to remember as you can approximate many problems as uniform density spheres.

(d) As the Sun contracted to its present radius, the virial theorem states that half of the total gravitational potential energy  $(U_g/2)$  was converted into kinetic energy and half radiated away. Assuming it is ok to approximate the Sun as a uniform density sphere, at the present luminosity of the Sun, how long would it take the Sun to radiate that energy away (answer in years)? Can this account for the 4.6 Gyr lifetime of the Sun?