

Example Neutral (H or H₂) Collision Rates

Table 4.2. Rate Coefficients (cm³s⁻¹) for Deexcitation by H and H₂

| Temperature T (°K) | 10° | 30° | 100° | 300° | 1000° |
|-----------------------------------------------------|------|------|------|------|-------|
| $10^{11} \times \gamma_{ki}$ for H–H | 0.23 | 3.0 | 9.5 | 16 | 25 |
| $10^{10} \times \gamma_{kj}$ for H-C ⁺ | 6.9 | 7.4 | 8.0 | 8.4 | 9.7 |
| $10^{12} \times \gamma_{20}$ for H-H ₂ | 0.96 | 1.37 | 3.0 | 9.1 | 42 |
| $10^{11} \times \gamma_{10}$ for H ₂ -CO | 1.8 | 3.2 | 3.7 | | |

Spitzer - Physical Processes in the ISM

| Table 4.1. | Collisi | | | Excitation by Electrons | | |
|-----------------------|---------|-------------------------------|-------------------------------|-------------------------|---------------|--------------------------|
| Number of p electrons | ion | Lower | Upper | $E_{jk}(eV)$ | $\Omega(j,k)$ | $\sum_{j}A_{kj}(s^{-1})$ |
| 1,5 | CII | ² P _{1/2} | ² P _{3/2} | 0.0079 | 1.33 | 2.4×10 ⁻⁶ |
| | Ne II | $^{2}P_{3/2}$ | | 0.097 | 0.37 | 8.6×10^{-3} |
| | Si II | $^{2}P_{1/2}$ | | 0.036 | 7.7 | 2.1×10^{-4} |
| 2 | NII | $^{3}P_{0}$ — | | 0.0061 | 0.41 | 2.1×10^{-6} |
| | | $^{3}P_{0}$ — | | 0.0163 | 0.28 | 7.5×10^{-6} |
| | | $^{3}P_{1}$ — | $^{3}P_{2}$ | 0.0102 | 1.38 | 7.5×10^{-6} |
| | | ³ P — | | 1.90 | 2.99 | 4.0×10^{-3} |
| | | ³ P — | | 4.05 | 0.36 | 1.1 |
| | OIII | $^{3}P_{0}$ — | | 0.014 | 0.39 | 2.6×10^{-5} |
| | | ${}^{3}P_{0}$ — | | 0.038 | 0.21 | 9.8×10^{-5} |
| | | $^{3}P_{1}$ — | | 0.024 | 0.95 | 9.8×10^{-5} |
| | | ³ P — | | 2.51 | 2.50 | 2.8×10^{-2} |
| 3 | OII | ³ P — | | 5.35 | 0.30 | 1.8 |
| | OII | 4S _{3/2} | | 3.32 | 0.88 | 4.2×10^{-5} |
| Physical | | $^{4}S_{3/2}$ | | 3.32 | 0.59 | 1.8×10^{-4} |
| es in the ISM | | $^{2}D_{3/2}$ | -D5/2 | 0.0025 | 1.16 | 4.2×10^{-5} |

